

# Study Links Autism to Vaccines Made With Cells From Aborted Babies

by Katie Doan | LifeNews.com | 9/9/14 4:20 PM

A new study published in the September 2014 volume of the Journal of Public Health and Epidemiology reveals a significant correlation between autism disorder (AD) and MMR, Varicella (chickenpox) and Hepatitis-A vaccines.

Using statistical analysis and data from the US Government, UK, Denmark and Western Australia, scientists at Sound Choice Pharmaceutical Institute (SCPI) found that increases in autistic disorder correspond with the introduction of vaccines using human fetal cell lines and retroviral contaminants.



Even more alarming, Dr Theresa Deisher, lead scientist and SCPI founder noted that, “Not only are the human fetal contaminated vaccines associated with autistic disorder throughout the world, but also with epidemic childhood leukemia and lymphomas.”

Their study comes on the heels of recent breaking news that the CDC deliberately withheld evidence of the significant increase in autism among African-American boys who were vaccinated prior to 36 months of age. (See:

[www.examiner.com/article/whistleblower-reveals-](http://www.examiner.com/article/whistleblower-reveals-cdc-cover-up-linking-mmr-vaccine-to-autism)

[cdc-cover-up-linking-mmr-vaccine-to-autism](http://www.examiner.com/article/whistleblower-reveals-cdc-cover-up-linking-mmr-vaccine-to-autism) )

So it should come as no surprise that the FDA has known for decades about the dangers of insertional mutagenesis by using the human fetal cell lines and yet, they chose to ignore it. Instead of conducting safety studies they regulated the amount of human DNA that could be present in a vaccine to no greater than 10ng. ([www.fda.gov/ohrms/dockets/ac/05/slides/5-4188S1\\_4draft.ppt](http://www.fda.gov/ohrms/dockets/ac/05/slides/5-4188S1_4draft.ppt) )

Unfortunately, Dr. Deisher’s team discovered that the fetal DNA levels ranged anywhere from 142ng – 2000ng per dose, way beyond the so-called “safe” level.

“There are a large number of publications about the presence of HERV (human endogenous retrovirus – the only re-activatable endogenous retrovirus) and its association with childhood lymphoma,” noted Dr Deisher. “The MMR II and chickenpox vaccines and indeed all vaccines that were propagated or manufactured using the fetal cell line WI-38 are contaminated with this retrovirus. And both parents and physicians have a right to know this!”

Certainly these discoveries by SCPI should generate an immediate investigation by FDA officials, if not an

outright ban on the use of aborted fetal cell lines as substrates for vaccine production. There are numerous other non-human FDA-approved cell lines that can and should be used.

Dr Deisher's study is available on the Academic Journals website at:

[www.ms.academicjournals.org/article/article1409245960\\_Deisher%20et%20al.pdf](http://www.ms.academicjournals.org/article/article1409245960_Deisher%20et%20al.pdf)

or on their website at [www.soundchoice.org/scpiJournalPubHealthEpidem092014.pdf](http://www.soundchoice.org/scpiJournalPubHealthEpidem092014.pdf)